

MASS-RADIUS RELATION FOR MAGNETIC WHITE DWARFS

IN-SAENG SUH¹ AND G. J. MATHEWS²

Center for Astrophysics, Department of Physics, University of Notre Dame, Notre Dame, Indiana 46556, USA

¹isuh@cygnus.phys.nd.edu; ²gmathews@bootes.phys.nd.edu*To appear in ApJ (Feb 20, 2000), vol. 530 (astro-ph/9906239)*

ABSTRACT

Recently, several white dwarfs with very strong surface magnetic fields have been observed. In this paper we explore the possibility that such stars could have sufficiently strong internal fields to alter their structure. We obtain a revised white dwarf mass-radius relation in the presence of strong internal magnetic fields. We first derive the equation of state for a fully degenerate ideal electron gas in a magnetic field using an Euler-MacLaurin expansion. We use this to obtain the mass-radius relation for magnetic ⁴He, ¹²C, and ⁵⁶Fe white dwarfs of uniform composition.

Subject headings: stars: white dwarfs - stars: magnetic fields - stars: interiors

1. INTRODUCTION

A number of white dwarfs with strong magnetic fields have been discovered (Kemp et al. 1970; Putney 1995; Schmidt & Smith 1995; Reimers et al. 1996) and extensively studied (Jordan 1992; Angel 1978; Channugam 1992 and references therein). Surface magnetic fields ranging from about 10⁵ G to 10⁹ G have been detected in about 50 (2%) of the ≈ 2100 known white dwarfs (Jordan 1997 and references therein). As relics of stellar interiors, the study of the magnetic fields in and around degenerate stars should give important information on the role such fields play in star formation and stellar evolution. However, the origin and evolution of stellar magnetic fields remains obscure.

As early as Ginzburg (1964) and Woltjer (1964) it was proposed that the magnetic flux ($\Phi_B \sim BR^2$) of a star is conserved during its evolution and subsequent collapse to form a remnant white dwarf or neutron star. A main sequence star with radius on the order of $R \sim 10^{11}$ cm and surface magnetic field $B \sim 10 - 10^4$ G [magnetic A-type stars have typical surface fields $\lesssim 10^4$ G (Shapiro & Teukolsky 1983)] would thus collapse to form a white dwarf with $R \sim 10^9$ cm and $B \sim 10^5 - 10^8$ G, or a neutron star with $R \sim 10^6$ cm and $B \sim 10^{11} - 10^{14}$ G. Indeed, shortly after their discovery (Hewish et al. 1968) pulsars were identified as rotating neutron stars (Gold 1968) with magnetic fields $B \sim 10^{11} - 10^{13}$ G consistent with magnetic field amplification by flux conservation. In addition, neutron stars with surface magnetic fields exceeding 10^{14} G [so called magnetars] have been recently suggested as the source of soft gamma-ray repeaters (Duncan & Thompson 1992; Thompson & Duncan 1995).

Moreover, the surface magnetic field of a star does not necessarily reflect the internal field (Ruderman 1980). For example, the toroidal fields below the surface of the Sun are at least on the order of $\sim 10^2$ to $\sim 10^4$ times stronger than the average surface dipole field strength of ~ 1 G (Galloway, Proctor, & Weiss 1977). Furthermore, at the region of the convective zone, the strength of small scale magnetic fields could reach a value as high as 7×10^4 G (Chauhan, Pandey & Pandey 1999, Pulido 1998). This would correspond to an interior field strength on the order of $\sim 10^9$ to $\sim 10^{13}$ G in a white dwarf, or $\sim 10^{15}$ to $\sim 10^{18}$ G in a neutron star. [Condensed objects of size R and mass M have an upper limit to their field strengths

of $B \lesssim MR^2(8\pi G)^{1/2}$. For neutron stars with $R \approx 10$ km and $M \approx M_\odot$, the limit is $B \lesssim \sim 10^{18}$ G (Lerche & Schramm 1977).]

Indeed, the existence of white dwarfs with interior magnetic fields as strong as $\sim 4 \times 10^{13}$ G is not ruled out with the present uncertainties in the mass-radius relation (Shapiro & Teukolsky 1983). The present high upper limit on the strength of internal fields in white dwarfs is obtained by simply setting the magnetic pressure equal to the internal pressure of the star. However, white dwarfs with internal fields at or around this strength could be constrained (Mestel 1965) by a perceptibly different mass-radius relation.

Although white dwarfs in binaries with well determined masses do not appear to have surface magnetic fields larger than $\sim 10^5$ G, internal fields of order 10^{12} G could be well hidden below the surface (Angel 1978). Newly discovered magnetic degenerate stars, especially those with surface field strengths near the range of $B \sim 10^9$ G, always show strong circularly and/or linearly polarized spectral energy distributions (Schmidt et al. 1999). Moreover, these stars reveal unique spectral features (Engelhardt & Bues 1994) due to quasi-Landau resonances in extremely high magnetic fields of $> 10^9$ G.

In this work, we explicitly compute the mass-radius relation of white dwarfs with internal magnetic fields. Previously, Ostriker & Hartwick (1968) have estimated effects of interior magnetic fields by considering a correction in terms of the ratio of magnetic to gravitational energy. They showed that a relatively small ratio of magnetic to gravitational energy would be sufficient to explain an observational discrepancy in the classical mass-radius relation for Sirius B. However, if white dwarfs could indeed have central magnetic fields as strong as $4.4 \times 10^{11} - 4.4 \times 10^{13}$ G, the revised mass-radius relation must be explicitly determined by taking the magnetic field into account in the equation of state. The present work thus expands upon that earlier study by explicitly computing the equation of state for a completely degenerate, noninteracting electron gas in a magnetic field. This equation of state is then applied to the Tolman-Oppenheimer-Volkoff (TOV) equation of stellar hydrostatic equilibrium.

The equation of state in a magnetic field should reduce to a normal equation of state in the absence of a magnetic field. Therefore, we use an Euler-MacLaurin expansion (Kernan, Starkman, & Vachaspati 1996) of the thermodynamic variables to recover the weak field limit. In integrating the TOV equation

we simply follow the procedure of Hamada & Salpeter (1961) for degenerate matter of uniform composition of ${}^4\text{He}$, ${}^{12}\text{C}$, or ${}^{56}\text{Fe}$. Although ${}^4\text{He}$ and ${}^{12}\text{C}$ white dwarfs are expected to have a similar (though not identical) mass-radius relation, we explicitly consider each for completeness.

2. EQUATION OF STATE FOR AN ELECTRON GAS IN A MAGNETIC FIELD

The properties of an electron in an external magnetic field have been studied extensively (Landau & Lifshitz 1938; Johnson & Lippmann 1949; Canuto & Chiu 1968; Schwinger 1988). In brief, the energy states of an electron in a magnetic field are quantized and its properties are modified accordingly. In order to investigate these effects, we must first solve the Dirac equation in an external, static, and homogeneous magnetic field. We make the convenient choice of gauge for the vector potential in which a uniform magnetic field B lies along the z -axis. We then obtain the electron wavefunctions and energy dispersion in a magnetic field (Johnson & Lippmann 1949). The dispersion relation for an electron propagating through a magnetic field is

$$E_n = [p^2 c^2 + m^2 c^4 + 2\hbar c e B n]^{1/2}, \quad (1)$$

where $n = j + \frac{1}{2} + s_z$, ($n = 0, 1, \dots$), j is the principal quantum number of the Landau level, $s_z = \pm 1/2$ is the electron spin, e is the electron charge, c is the speed of light, $\hbar$ is Planck's constant, $p \equiv p_z$ is the electron momentum along the z -axis, and m is the rest mass of the electron. In Eq. (1) we ignore the anomalous magnetic moment for an electron.

The main modification of an electron in a magnetic field comes from the available density of states for the electrons (Landau & Lifshitz 1938). The electron state density in the absence of a magnetic field,

$$\frac{2}{h^3} \int \frac{d^3 \vec{p}}{(2\pi)^3},$$

is replaced with

$$\frac{2}{h^2 c} \sum_{n=0}^{\infty} [2 - \delta_{n0}] \int \frac{eB}{(2\pi)^2} dp_z$$

in a magnetic field. The symbol δ_{n0} is the Kronecker delta defined by $\delta_{n0} = 1$ for $n = 0$, or $\delta_{n0} = 0$ for $n \neq 0$. This modification affects the thermodynamic variables for the electron gas.

An isolated white dwarf ultimately cools to zero temperature (fully degenerate) and the degeneracy pressure supports these stars against further gravitational collapse. For the most part, this pressure can be described as an ideal (noninteracting) electron gas plus small corrections.

Let us consider a gas of electrons at zero temperature in a magnetic field. From Eq. (1) we can define the Fermi energy E_F for an arbitrary Landau level n as

$$E_F^2 \equiv m^2 c^4 + p_F^2 c^2 + 2\hbar c e B n. \quad (2)$$

Here p_F denotes the Fermi momentum. The number density of electrons in a magnetic field is then given by

$$n_e = 2 \frac{\gamma}{(2\pi)^2} \left(\frac{mc}{\hbar} \right)^3 \zeta(\epsilon_f, n), \quad (3)$$

where

$$\zeta(\epsilon_f, n) = \sum_{n=0}^{n_f} [2 - \delta_{n0}] \sqrt{\epsilon_f^2 - (1 + 2\gamma n)}.$$

In the above, $\epsilon_f \equiv E_F/mc^2$ and $\gamma = B/B_c$ where $B_c = m^2 c^3 / |e| \hbar = 4.414 \times 10^{13}$ G is the critical magnetic field at which quantized cyclotron states begin to exist. The maximum Landau level n_f for a given Fermi energy ϵ_f and magnetic field strength γ is given by

$$n_f \equiv \frac{\epsilon_f^2 - 1}{2\gamma} \geq n.$$

The pressure of an ideal electron gas in a magnetic field is then

$$P_e = 2 \frac{\gamma}{4\pi^2} mc^2 \left(\frac{mc}{\hbar} \right)^3 \Phi(\epsilon_f, n), \quad (4)$$

where

$$\begin{aligned} \Phi(\epsilon_f, n) = & \frac{1}{2} \sum_{n=0}^{n_f} [2 - \delta_{n0}] \left[\epsilon_f \sqrt{\epsilon_f^2 - (1 + 2\gamma n)} \right. \\ & \left. - (1 + 2\gamma n) \ln \left(\frac{\epsilon_f + \sqrt{\epsilon_f^2 - (1 + 2\gamma n)}}{\sqrt{1 + 2\gamma n}} \right) \right]. \end{aligned}$$

Similarly, the energy density is

$$\mathcal{E}(\epsilon_f, n) = 2 \frac{\gamma}{4\pi^2} mc^2 \left(\frac{mc}{\hbar} \right)^3 \chi(\epsilon_f, n), \quad (5)$$

where

$$\begin{aligned} \chi(\epsilon_f, n) = & \frac{1}{2} \sum_{n=0}^{n_f} [2 - \delta_{n0}] \left[\epsilon_f \sqrt{\epsilon_f^2 - (1 + 2\gamma n)} \right. \\ & \left. + (1 + 2\gamma n) \ln \left(\frac{\epsilon_f + \sqrt{\epsilon_f^2 - (1 + 2\gamma n)}}{\sqrt{1 + 2\gamma n}} \right) \right]. \end{aligned}$$

From these, we obtain the energy per electron

$$E_e(\epsilon_f, n) = mc^2 \frac{\chi(\epsilon_f, n)}{\zeta(\epsilon_f, n)}. \quad (6)$$

In order to recover the usual equation of state in the absence of a magnetic field, we utilize an Euler-MacLaurin expansion of Eqs. (3) - (6) in the weak field limit. Then, the number density is given by

$$n_e \simeq \frac{1}{3\pi^2} \left(\frac{mc}{\hbar} \right)^3 \zeta(x), \quad (7)$$

where,

$$\zeta(x) = x^3 + \gamma^2 \frac{1}{4x} + \mathcal{O}(\gamma^4) + \dots,$$

and $x \equiv p_F/mc$ is the relativity parameter. Note that the electron number density increases as the magnetic field increases for a given x .

The pressure becomes

$$P_e \simeq \frac{1}{24\pi^2} mc^2 \left(\frac{mc}{\hbar} \right)^3 \Phi(x), \quad (8)$$

where

$$\Phi(x) = \Phi_0(x) + \gamma^2 \Phi_B(x) + \mathcal{O}(\gamma^4) + \dots,$$

$$\Phi_0(x) = x\sqrt{x^2+1}(2x^2-3) + 3\ln(x + \sqrt{x^2+1}),$$

$$\Phi_B(x) = \frac{\sqrt{x^2+1}}{x} + 2\ln(x + \sqrt{x^2+1}) - \left(1 + \frac{1}{x(x + \sqrt{x^2+1})}\right).$$

Note also that for a physically reasonable value of x the pressure always increases as γ increases.

The energy density can also be written

$$\mathcal{E}(x) \simeq mc^2 \left(\frac{mc}{\hbar}\right)^3 \chi(x), \quad (9)$$

where

$$\begin{aligned} \chi(x) &= \chi_0(x) + \gamma^2 \chi_B(x) + \mathcal{O}(\gamma^4) + \dots, \\ \chi_0(x) &= \frac{1}{8\pi^2} \left[x(2x^2+1)\sqrt{x^2+1} - \ln(x + \sqrt{x^2+1}) \right], \\ \chi_B(x) &= \frac{1}{24\pi^2} \left[1 + \frac{\sqrt{x^2+1}}{x} + \frac{1}{x(x + \sqrt{x^2+1})} - 2\ln(x + \sqrt{x^2+1}) \right]. \end{aligned}$$

Finally, the energy per electron is given by

$$E_e(x) \simeq \frac{3}{8} mc^2 \frac{\chi(x)}{\zeta(x)}. \quad (10)$$

Here we can see explicitly that as γ goes to zero, Eqs. (7) - (10) recover exactly the usual equation of state in the absence of a magnetic field.

3. MASS-RADIUS RELATION OF MAGNETIC WHITE DWARFS

The mass-radius relation of white dwarfs was first determined by Chandrasekhar (1939). Later Hamada & Salpeter (1961) obtained numerical models for various core compositions by considering a fully degenerate configuration at zero temperature. The theoretical relationship between the mass and radius of a white dwarf is important for the interpretation of observational results (see Koester & Chanmugam 1990 for a review). There are several recent studies and observations on the mass-radius relation of non-magnetic white dwarfs (Wood 1990; Vauclair, Schmidt, & Koester 1997; Vennes, Fontaine, & Brassard 1995; Provencal et al. 1998).

In order to obtain the mass-radius relation for magnetic white dwarfs, we use Eqs. (7) - (10) for $\gamma \leq 1$ ($B \leq 4.4 \times 10^{13}$ G) and carry out stellar integrations for a uniform composition of ${}^4\text{He}$, ${}^{12}\text{C}$, and ${}^{56}\text{Fe}$ as an illustrative model. In this uniform model (Hamada & Salpeter 1968; Fushiki et al. 1992; R  gnvaldsson et al. 1993) the total energy E of the plasma consists of a nearly uniform distribution of degenerate electrons with embedded ions,

$$E = E_e + E_C, \quad (11)$$

where the first term is the energy of a uniform gas of free electrons and E_C corrects for the classical Coulomb energy. Although the noninteracting electron gas accounts for the dominant contribution to the equation of state at high density, the classical Coulomb correction is significant. Other corrections,

such as the Thomas-Fermi, exchange, and correlation corrections give only a very small change in the mass-radius relation of white dwarfs. Actually the Thomas-Fermi correction in a strong magnetic field is important at low density (see R  gnvaldsson et al. 1993 and references therein), but as a whole it gives only a minor effect on the mass-radius relation. Hence, we ignore these minor effects for the present work.

Magnetic fields should not alter the spherical symmetry of the constituent atoms. Indeed, it is, perhaps, a remarkable fact that magnetic fields do not destroy even the approximate spherical symmetry of heavy atoms within the relevant range (R  gnvaldsson et al. 1993). Thus, we can use the ordinary electrostatic energy (Coulomb energy) per charge,

$$E_C/Z = -\frac{9}{5} Z^{2/3} \text{Ry} \frac{1}{r_e}, \quad (12)$$

where $\text{Ry} = \frac{1}{2} \alpha^2 m_e c^2$ is the Rydberg energy and from Eq. (7),

$$r_e = \left(\frac{3\pi}{8\gamma\zeta(x)} \right)^{1/3} \alpha.$$

The corresponding pressure is

$$P_C = -mc^2 \left(\frac{mc}{\hbar}\right)^3 \frac{9}{4\pi} \frac{Z^{2/3} \alpha^5}{10} \frac{1}{r_e^4}. \quad (13)$$

In this model, the total pressure is then given by

$$P = P_e + P_C. \quad (14)$$

Thus, since P_C is negative, the equation of state would lead to negative pressures at low density (Salpeter 1961).

In integrating the equations of hydrostatic equilibrium, we follow the classical procedure of Hamada & Salpeter (1961). Figures 1, 3, and 5 show the mass-radius relation of ${}^4\text{He}$, ${}^{12}\text{C}$, and ${}^{56}\text{Fe}$ white dwarfs for a given magnetic field strength. Figures 2, 4, and 6 show the relation between mass and central density of these white dwarfs for a given magnetic field strength. Here it can be seen that our results approach the Hamada & Salpeter (1961) results as the magnetic field strength decreases. For high central field strengths $\gamma \simeq 0.01 - 1$ [$B \simeq 4.4 \times (10^{11} - 10^{13})$ G], both the mass and radius of magnetic white dwarfs increase compared to non-magnetic white dwarfs of the same central density. For instance, for $\gamma \approx 0.8$ carbon white dwarfs, the radius increases by about 30% for $M \approx 1 M_\odot$. Similarly for $R \approx 0.01 R_\odot$ the mass also increases by about 25%. These results are approximately consistent with Ostriker & Hartwick (1968), i.e. the radius increases while the central density ρ_c decreases as the magnetic field increases for fixed M . As expected, for $B \lesssim 10^{10}$ G, internal magnetic fields do not affect the white-dwarf mass-radius relation.




FIG. 1 Relation between the mass M and radius R of a ${}^4\text{He}$ magnetic white dwarf for the indicated magnetic-field strengths. The solid line denotes the Hamada & Salpeter model for non-magnetic white dwarfs ($\gamma = 0$). The dashed lines are magnetic white dwarfs.




FIG. 2 Relation between the central density ρ_c (in g/cm^3) and mass M for ${}^4\text{He}$ magnetic white dwarfs. The solid line denotes the Hamada-Salpeter model for non-magnetic white dwarfs ($\gamma = 0$). The dashed lines are for magnetic white dwarfs.

A striking feature of these results is that white dwarfs with strong interior magnetic fields should be massive. This is simply because a star becomes unbound if the magnetic plus matter pressure force exceeds the gravitational force. For example, there is no stable solution with $M \lesssim 0.5M_\odot$ for $\gamma = 0.3$ carbon white dwarfs. This is consistent with recent observations that

on average magnetic white dwarfs have a higher mass than typical non-magnetic white dwarfs (see Table 1). Note, however, that for a field strength of $B \lesssim 1 \times 10^{13}$ G, magnetic fields only give a relatively small change on the mass-radius relation, that is, we can not distinguish between magnetic and non-magnetic white dwarfs for $\gamma \lesssim 0.2$.





4. DISCUSSION

In this work, we have calculated the equation of state for an electron gas in a magnetic field at zero temperature. For simplicity, we have assumed a uniform composition to obtain the relation between mass and radius for magnetic white dwarfs. For high internal magnetic fields $B \simeq 4.4 \times (10^{11} - 10^{13})$ G, ($\gamma \simeq 0.01 - 1$), the mass-radius relation is modified. Our results not only confirm the Ostriker & Hartwick (1968) result of increasing radius and decreasing central density ρ_c with increasing field, but also are consistent with the suggestion (Liebert 1988) that observed magnetic white dwarfs have masses which are on average larger than non-magnetic white dwarfs, implying more massive and younger progenitors.

The question remains, however, as to whether it is reasonable to consider such high internal field strengths for magnetic white dwarfs. First, one must assume that the magnetic fields are well hidden beneath the surface, while the surface fields are several orders of magnitude less. Second, assuming that flux is conserved during the collapse to a white dwarf, the progenitor of the white dwarfs must have had sufficiently large fields to produce the required white dwarf internal field strengths. Flux conservation implies that the central field strength of the progenitors is of order $\sim 10^8$ G, assuming $R \approx 1R_\odot$. This is reasonable. During star formation, the collapse of a typical interstellar cloud with radius ~ 0.1 pc, mass $\sim 1M_\odot$, and protostellar magnetic field of magnitude $\sim 3 \times 10^{-6}$ G would result in a field strength of $\sim 3 \times 10^8$ G in a solar type star formed from this material (Spitzer 1978). Although there is no evidence for main sequence stars with such field strengths, they are not ruled out by observations either (Shapiro & Teukolsky 1983).

5. COMPARISON WITH OBSERVATIONS

Figure 7 shows a comparison between our calculations and white dwarfs with known masses and radii from the HIPPARCOS survey (Vauclair, Schmidt, & Koester 1997; Provencal et al. 1998). Plotted error bars are the quoted $\pm 1\sigma$ observational uncertainties. Strong hidden interior magnetic fields would be expected to manifest themselves by a preponderance of stars with large masses and radii. Most of the data, however, are within 2σ of the non-magnetic theoretical curves. A puzzling feature however (Provencal et al. 1998) is that some of the best determined data points (e.g., EG 50 and Procyon B) can not be fit without postulating an iron composition, something which seems unlikely from a stellar evolution standpoint. Furthermore, although the evidence is not compelling, there are at least two well determined stars [i.e., GD 140 and Sirius B (Provencal et al. 1998)] as well as some field stars with $M \gtrsim 0.6M_\odot$ and $R \gtrsim 0.012R_\odot$ which may be better fit if strong internal magnetic fields are assumed. However, Sirius B and some of the field stars can also be accommodated by atmospheric models (Wood 1990).

FIG. 3 Same as Fig. 1, but for ^{12}C .FIG. 5 Same as Fig. 1, but for ^{56}Fe .FIG. 4 Same as Fig. 2, but for ^{12}C .FIG. 6 Same as Fig. 2, but for ^{56}Fe .




FIG. 6 A comparison of white dwarfs with known masses and radii from the HIPPARCOS survey at the 1σ level (Vauclair, Schmidt, & Koester 1997; Provencal et al. 1998) with our calculations from Figs. 1, 3, and 5. The upper curves are for ${}^4\text{He}$ and ${}^{12}\text{C}$ white dwarfs. The lower curves are for ${}^{56}\text{Fe}$ white dwarfs.

Some typical magnetic white dwarfs which have reported masses M and surface magnetic fields B_s are summarized in Table 1. This table shows that magnetic white dwarfs are typically more massive than non-magnetic white dwarfs (on average $M \sim 0.6M_\odot$). Among them RE J0317-858 ($B_s > 10^8$ G) is approaching the Chandrasekhar mass limit (Barstow et al.

1995).

Not listed in Table 1, however, are the two known magnetic white dwarfs with the strongest surface magnetic field, GD 229 ($B_s \gtrsim 10^9$ G) and PG 1031+234 ($B_s \approx (0.5-1) \times 10^9$ G). Unfortunately, there are no reported masses or radii for those magnetic white dwarfs. If their masses and radii could be measured, then our magnetic field model might be tested.

Perhaps the most interesting object in Table 1 is LB 11146 (PG 0945+245). This is an unresolved binary system consisting of two degenerate stars (Liebert et al. 1993): one component is a normal DA white dwarf ($M = 0.91 \pm 0.07 M_\odot$) with no detectable magnetic field; the other has a strong magnetic field ($B_s > 3 \times 10^8$ G) and a mass in the range $0.76 \leq M/M_\odot \lesssim 1.0$. Thus, assuming for illustration a strong interior field of $\gamma \approx 0.5$, and that this is a typical equal mass system with $M = 0.9M_\odot$ each, then the radius of the magnetic component would be larger by about 10% than the normal star. Alternatively, if they have similar radii of $R = 0.01 R_\odot$, the magnetic star would have a heavier mass by about 12% than the normal one for $\gamma \approx 0.5$.

Regarding the radius of magnetic white dwarfs, Greenstein & Oke (1982) have reported radii of $R \sim 0.0066 R_\odot$ for Grw +70°8247 ($B_s \approx 3.2 \times 10^8$ G) and $R \sim 0.01 R_\odot$ for Feige 7 ($B_s \approx 3.5 \times 10^7$ G) from the interpretation of their spectra. Assuming an interior field which is about 10^5 times stronger than the surface field, then we would deduce $\gamma \approx 0.8$ and $M \sim 1.1-1.2 M_\odot$ for Grw +70°8247, consistent with the expectation of high masses for magnetic white dwarfs. Similarly, we would deduce $\gamma \approx 0.075$ and $M \sim 0.8 M_\odot$ for Feige 7. This star however would be nearly indistinguishable from a non-magnetic white dwarf.

This work supported in part by NSF Grant-97-22086 and DOE Nuclear Theory Grant DE-FG02-95ER40934.

REFERENCES

- Angel, J. R. P. 1978, *ARA&A* 16, 487
 Barstow, M. A., Jordan, S., O'Donoghue, D., Burleigh, M. R., Napiwotzki, R., & Harrop-Allin, M. K. 1995, *MNRAS* 277, 971
 Canuto, V. & Chiu, H.-Y. 1968, *Phys. Rev.* 173, 1210
 Chandrasekhar, S. 1939, *An Introduction to the Study of Stellar Structure*, (Chicago: Univ. Chicago Press)
 Chanmugam, G. 1992, *ARA&A* 30, 143
 Chauhan, B. C., Pandey, U. C., and Dev, S. 1999 *Phys. Rev. D* 59, 083002
 Duncan, R. C. & Thompson, C. 1992, *ApJ* 392, L9
 Engelhardt, D. & Bues, I. 1994, in *White Dwarfs*, ed. D. Koester & K. Werner (Berlin: Springer) 123
 Ferrario, L., Vennes, S., & Wickramasinghe, D. T. 1998, *MNRAS* 299, L1
 Fushiki, I., Gudmundsson, E. H., Pethick, C. J., & Yngvason, J. 1992, *Ann. Phys.* 216, 29
 Galloway, D. J., Proctor, M. R. E., & Weiss, N. O. 1977, *Nature* 266, 686
 Ginzberg, V. L. 1964, *Sov. Phys. Dokl.* 9, 329
 Gold, T. 1968, *Nature* 218, 731
 Greenstein, J. L. & Oke, J. B. 1982, *ApJ* 252, 285
 Hamada, T. & Salpeter, E. E. 1961, *ApJ* 134, 683
 Hewish, A., Bell, S. J., Pilkington, J. D., Scott, P. F., & Collins, R. A. 1968, *Nature* 217, 709
 Johnson, M. H. & Lippmann, B. A. 1949, *Phys. Rev.* 76, 828
 Jordan, S. 1992, *A & A* 265, 570
 Jordan, S. 1997, in *White Dwarfs*, ed. J. Isern, M. Hernanz, & E. Garcia-Berro, 397
 Kemp, J. C., Swedlund, J. B., Landstreet, J. D., & Angel, J. R. P. 1970, *ApJ* 161, L77
 Kernan, P. J., Starkman, G. D., & Vachaspati, T. 1996, *Phys. Rev. D* 54, 7207
 Koester, D. & Chanmugam, G. 1990, *Rep. Prog. Phys.* 53, 837
 Landau, L. D. & Lifshitz, E. M. 1938, *Statistical Mechanics* (Clarendon: Oxford)
 Lerche, I. & Schramm, D. N. 1977, *ApJ* 216, 881
 Liebert, J. 1988, *PASP* 100, 1302
 Liebert, J., Bergeron, P., Schmidt, G. D., & Saffer, R. A. 1993, *ApJ* 418, 426
 Moran, C., Marsh, T. R., & Dhillon, V. S. 1998, *MNRAS* 299, 218
 Mestel, L. 1965, *Stars and Stellar Systems*, ed. L. H. Aller & D. B. McLachlin, (Chicago: Univ. Chicago Press) 297
 Ostriker, J. P. & Hartwick, F. D. A. 1968, *ApJ* 153, 797
 Provencal, J. L., Shipman, H. L., Hog, E., & Thejll, P. 1998, *ApJ* 494, 759
 Pulido, J. 1998, *Phys. Rev. D* 57, 7108
 Putney, A. 1995, *ApJ* 451, L67
 Reimers, D., Jordan, S., Koester, D., Bade, N., Köhler, Th. & Wisotzki, L. 1996, *A & A* 311, 572
 Rögnvaldsson, Ö. E., Fushiki, I., Gudmundsson, E. H., Pethick, C. J., & Yngvason, J. 1993, *ApJ* 416, 276
 Ruderman, M. 1980, *J. de Physique suppl.* 41, C2-125
 Salpeter, E. E. 1961, *ApJ* 134, 669
 Schmidt, G. D., Bergeron, P., Liebert, J., & Saffer, R. 1992, *ApJ* 394, 603
 Schmidt, G. D., Liebert, J., Harris, H. C., Dahn, C. C., & Leggett, S. K. 1999, *ApJ* 512, 916
 Schmidt, G. D. & Smith, P. S. 1995, *ApJ* 448, 305
 Schwinger, J. 1988, *Particles, Sources and Fields* (Redwood City: Addison-Wesley)
 Shapiro, S. L. & Teukolsky, A. A. 1983, *Black Holes, White Dwarfs, and Neutron Stars* (New York: Wiley-Interscience)
 Spitzer, L. 1978, *Physical Processes in the Interstellar Medium* (New York, Wiley-Interscience)
 Thompson, C. & Duncan, R. C. 1995, *MNRAS* 275, 255
 Vauclair, G., Schmidt, H., Koester, D., & Allard, N. 1997, *A & A* 325, 1055
 Vennes, S., Fontaine, G., & Brassard, P. 1995, *A & A* 296, 117
 Vennes, S., Thejll, P. A., Galvan, R. G., & Dupuis, J. 1997, *ApJ* 480, 714
 Woltjer, L. 1964, *ApJ* 140, 1309
 Wood, M. A. 1990, Ph.D. thesis, Univ. of Texas at Austin

TABLE 1
MASS AND SURFACE MAGNETIC FIELD STRENGTH IN SOME TYPICAL MAGNETIC WHITE DWARFS

	$M(M_{\odot})$	$B_s(10^6 \text{ G})$	Ref.
PG 2329+267	~ 0.9	2.3	Moran et al. 1998
LB 11146B	0.76 - 1.0	> 300	Liebert et al. 1993
Grw +70°8247	> 1.0	320	Greenstein & Oke 1990
1RXS J0823.6-2525	1.2	~ 3	Ferrario et al. 1998
PG 0136+251	1.28	1.3?	Vennes et al. 1997
PG 1658+441	1.31	3.5?	Schmidt et al. 1992
RE J0317-858	1.35	660	Barstow et al. 1995